

Polar bears finding it harder to catch enough seals to meet energy demands

Study reveals high metabolic rates, meaning polar bears need a lot of fat-rich prey, but more than half of those studied were running an energy deficit

Date: February 1, 2018

Source: University of California - Santa Cruz

Summary: A new study finds polar bears in the wild have higher metabolic rates than previously thought, and as climate change alters their environment a growing number of bears are unable to catch enough prey to meet their energy needs.

FULL STORY



This is an adult female polar bear on the sea ice wearing a GPS satellite video-camera collar. GPS video-camera collars were applied to solitary adult female polar bears for 8 to 12 days in April, 2014-2016. These collars enabled researchers to understand the movements, behaviors, and foraging success of polar bears on the sea ice.

Credit: Anthony Pagano, USGS

A new study finds polar bears in the wild have higher metabolic rates than previously thought, and as climate change alters their environment a growing number of bears are unable to catch enough prey to meet their energy needs.

The study, published February 2 in *Science*, reveals the physiological mechanisms behind observed declines in polar bear populations, said first author Anthony Pagano, a Ph.D. candidate at UC Santa Cruz.

"We've been documenting declines in polar bear survival rates, body condition, and population numbers over the past decade," he said. "This study identifies the mechanisms that are driving those declines by looking at the actual energy needs of polar bears and how often they're able to catch seals."

Pagano, who is also a wildlife biologist with the U.S. Geological Survey (USGS), conducted the study as part of his Ph.D. thesis research at UC Santa Cruz, where he has been working with coauthors Terrie Williams and Daniel Costa, both professors of ecology and evolutionary biology.

The researchers monitored the behavior, hunting success, and metabolic rates of adult female polar bears without cubs as they hunted for prey on the sea ice of the Beaufort Sea in the spring. High-tech collars on the bears recorded video, locations, and activity levels over a period of eight to 11 days, while metabolic

tracers enabled the team to determine how much energy the bears expended.

The field metabolic rates they measured averaged more than 50 percent higher than previous studies had predicted. Five of the nine bears in the study lost body mass, meaning they weren't catching enough fat-rich marine mammal prey to meet their energy demands.

"This was at the start of the period from April through July when polar bears catch most of their prey and put on most of the body fat they need to sustain them throughout the year," Pagano said.

Climate change is having dramatic effects on the Arctic sea ice, forcing polar bears to move greater distances and making it harder for them to catch prey. In the Beaufort Sea, sea ice starts to retreat away from the continental shelf in July, and most of the bears move north on the ice as it retreats. As the Arctic warms and more sea ice melts, the bears are having to move much greater distances than previously. This causes them to expend more energy during the summer, when they are fasting until the ice returns to the continental shelf in the fall.

In other areas, such as Hudson Bay, most bears move onto land when the sea ice retreats. There, Arctic warming means the sea ice is breaking up earlier in the summer and returning later in the fall, forcing bears to spend more time on land.

"Either way, it's an issue of how much fat they can put on before the ice starts to break up, and then how much energy are they having to expend," Pagano said.

Previous studies had tried to estimate polar bear metabolic rates and energy expenditures based on some assumptions about their behavior and physiology. For example, since polar bears are primarily "sit and wait" hunters, it was thought this would minimize their energy expenditure during hunting. Researchers also speculated that polar bears could lower their metabolic rate to save energy if they were not successful catching seals, Pagano said.

"We found that polar bears actually have much higher energy demands than predicted. They need to be catching a lot of seals," he said.

In the spring, polar bears are mostly preying on recently weaned ringed seals, which are more susceptible to being caught than adult seals. By the fall, the young seals are older and wiser, and polar bears are not able to catch as many. "It's thought that bears might catch a couple per month in the fall, compared to five to 10 per month in the spring and early summer," Pagano said.

USGS researchers have been studying polar bears in the Beaufort Sea area since the 1980s. Their most recent population estimate indicates the polar bear population has declined by about 40 percent over the past decade. It has been difficult, however, for researchers to study the fundamental biology and behavior of polar bears in this very remote and harsh environment, Pagano said.

"We now have the technology to learn how they are moving on the ice, their activity patterns, and their energy needs, so we can better understand the implications of these changes we are seeing in the sea ice," he said.

In addition to Pagano, Williams, and Costa, the coauthors of the paper include USGS researchers George Durner, Karyn Rode, Todd Atwood, and Elizabeth Peacock; Stephen Atkinson of Dugald, Manitoba; and Megan Owen at the San Diego Zoo Institute for Conservation Research. This work was supported by the USGS Changing Arctic Ecosystems Initiative.

Story Source:

Materials provided by **University of California - Santa Cruz**. Original written by Tim Stephens. *Note: Content may be edited for style and length.*

Journal Reference:

1. A. M. Pagano, G. M. Durner, K. D. Rode, T. C. Atwood, S. N. Atkinson, E. Peacock, D. P. Costa, M. A. Owen, T. M. Williams. **High-energy, high-fat lifestyle challenges an Arctic apex predator, the polar bear.** *Science*, 2018; 359 (6375): 568 DOI: 10.1126/science.aan8677

Cite This Page:

MLA

APA

Chicago

University of California - Santa Cruz. "Polar bears finding it harder to catch enough seals to meet energy demands: Study reveals high metabolic rates, meaning polar bears need a lot of fat-rich prey, but more than half of those studied were running an energy deficit." ScienceDaily. ScienceDaily, 1 February 2018. <www.sciencedaily.com/releases/2018/02/180201173314.htm>.

RELATED STORIES

Polar Bears' Declining Mercury Levels Likely Due to Climate-Related Shifts

June 14, 2017 — To understand how human activities are affecting the planet, scientists often study the health of animals in the wild. Now a new study finds that the levels of mercury in some polar bears are ... **read more »**



Polar Bears Are Swimming More as Sea Ice Retreats, Study Indicates

Apr. 20, 2016 — A study undertaken by scientists to understand swimming behavior in polar bears is showing an increase in this behavior related to changes in the amount and location of summer sea ice. The pattern of ... **read more »**



Polar Bears May Survive Ice Melt, With or Without Seals

Sep. 4, 2015 — As climate change accelerates ice melt in the Arctic, polar bears may find caribou and snow geese replacing seals as an important food source, shows a recent study. The research is based on new ... **read more »**

Scent Communication in Polar Bears Explored

Nov. 4, 2014 — Scientists have provided the first systematic examination of the social information polar bears may glean from scent left in the paw prints of other polar bears. The authors also suggest that scent ... **read more »**