

Polar bear diet changes as sea ice melts

Date: January 22, 2014

Source: American Museum of Natural History

Summary: At least some polar bears in the western Hudson Bay population are using flexible foraging strategies while on land, such as prey-switching and eating a mixed diet of plants and animals, as they survive in their rapidly changing environment, new research suggests.

FULL STORY



A polar bear, *Ursus maritimus*, eats a caribou.

Credit: Copyright American Museum of Natural History/R. Rockwell

A series of papers recently published by scientists at the American Museum of Natural History suggests that polar bears in the warming Arctic are turning to alternate food sources. As Arctic sea ice melts earlier and freezes later each year, polar bears have a limited amount of time to hunt their historically preferred prey -- ringed seal pups -- and must spend more time on land. The new research indicates that at least some polar bears in the western Hudson Bay population are using flexible foraging strategies while on land, such as prey-switching and eating a mixed diet of plants and animals, as they survive in their rapidly changing environment.

"There is little doubt that polar bears are very susceptible as global climate change continues to drastically alter the landscape of the northern polar regions," said Robert Rockwell, a research associate in the Museum's Department of Ornithology. "But we're finding that they might be more resilient than is commonly thought."

Polar bears are listed as a threatened species under the United States Endangered Species Act and are classified as "vulnerable" with declining populations on the International Union for Conservation of Nature and Natural Resources' Red List. Climate warming is reducing the availability of their ice habitat, especially

in the spring when polar bears gain most of their annual fat reserves by consuming seal pups before coming ashore for the summer. The new work, led by Rockwell and Linda Gormezano, a postdoctoral researcher in the Museum's Division of Vertebrate Zoology, examines how polar bears might compensate for energy deficits from decreasing seal-hunting opportunities.

In the first paper, published in spring 2013 in the journal *Polar Ecology*, the researchers provide, for the first time, data and video of polar bears pursuing, catching, and eating adult and juvenile lesser snow geese during mid-to-late summer, when the geese are replacing their primary flight feathers.

In the second paper, published in summer 2013 in the journal *Ecology and Evolution*, researchers used polar bear scat to show that the diet of at least some of the bears has shifted from what it was 40 years ago, before climate change was affecting the Hudson Bay lowlands. Today's polar bears are preying more on caribou as well as on snow geese and their eggs.

In the final paper in the series, published in December 2013 in the journal *BMC Ecology*, the researchers show that polar bears are, with a few exceptions, consuming a mixed diet of plants and animals. The predominance of local vegetation in collected scat suggests little movement among habitat types between feeding sessions, indicating that the polar bears are keeping energy expenditure down.

Taken together, the research indicates that during the ice-free period, polar bears are exhibiting flexible foraging behavior. This behavior likely derives from a shared genetic heritage with brown bears, from which polar bears separated about 600,000 years ago.

"For polar bear populations to persist, changes in their foraging will need to keep pace with climate-induced reduction of sea ice from which the bears typically hunt seals," Gormezano said. "Although different evolutionary pathways could enable such persistence, the ability to respond flexibly to environmental change, without requiring selective alterations to underlying genetic architecture, may be the most realistic alternative in light of the fast pace at which environmental changes are occurring. Our results suggest that some polar bears may possess this flexibility and thus may be able to cope with rapidly changing access to their historic food supply."

Funding for this work was provided by the Hudson Bay Project; the American Museum of Natural History; Churchill Northern Studies Centre; City University of New York; and Manitoba Conservation.

Story Source:

Materials provided by **American Museum of Natural History**. Note: Content may be edited for style and length.

Journal References:

1. D. T. Iles, S. L. Peterson, L. J. Gormezano, D. N. Koons, R. F. Rockwell. **Terrestrial predation by polar bears: not just a wild goose chase.** *Polar Biology*, 2013; 36 (9): 1373 DOI: 10.1007/s00300-013-1341-5
2. Linda J. Gormezano, Robert F. Rockwell. **What to eat now? Shifts in polar bear diet during the ice-free season in western Hudson Bay.** *Ecology and Evolution*, 2013; DOI: 10.1002/ece3.740
3. Linda J Gormezano, Robert F Rockwell. **Dietary composition and spatial patterns of polar bear foraging on land in western Hudson Bay.** *BMC Ecology*, 2013; 13 (1): 51 DOI: 10.1186/1472-6785-13-51

Cite This Page:

American Museum of Natural History. "Polar bear diet changes as sea ice melts." ScienceDaily. ScienceDaily, 22 January 2014. <www.sciencedaily.com/releases/2014/01/140122104025.htm>.

RELATED STORIES



Polar Bears Finding It Harder to Catch Enough Seals to Meet Energy Demands

Feb. 1, 2018 — A new study finds polar bears in the wild have higher metabolic rates than previously thought, and as climate change alters their environment a growing number of bears are unable to catch enough prey ... [read more »](#)

Polar Bears' Declining Mercury Levels Likely Due to Climate-Related Shifts

June 14, 2017 — To understand how human activities are affecting the planet, scientists often study the health of animals in the wild. Now a new study finds that the levels of mercury in some polar bears are ... [read more »](#)

Charcoaling Manure, Greening Neighborhoods in Chesapeake Bay Watershed

July 1, 2015 — Chesapeake Bay bears a heavy pollution burden from the growing metropolitan centers and vibrant agricultural activity in the watershed. Nitrogen and phosphorus draining from farm fields, livestock ... [read more »](#)



Algae Use Same Molecular Machinery as Land Plants to Respond to a Plant Hormone

Jan. 8, 2015 — Land-based plants -- including the fruits and vegetables in your kitchen -- produce and respond to hormones in order to survive. Scientists once believed that hormone signaling machinery only existed ... [read more »](#)